

On Newman-Penrose constants of stationary space-times

Xiaoning Wu^{a,1} and Yu Shang^{a,b,2}

a. Institute of Applied Mathematics,
Academy of Mathematics and System Science,
Chinese Academy of Sciences,
P.O.Box 2734, Beijing, China, 100080.
b. Graduate School of Chinese Academy of Science,
Beijing, China, 100080

Abstract

We consider the general asymptotic expression of stationary space-time. Using Killing equation, we reduce the dynamical freedom of Einstein equation to the ingoing gravitational wave Ψ_0 . The general form of this function can be got. With the help of asymptotically algebraic special condition, we prove that all Newman-Penrose constants vanish.

PACS number : 04.20.-q, 04.20.Ha

Keywords : Newman-Penrose constants, stationary solutions, algebraic special solutions.

1 Introduction

In a classical work by Newman and Penrose [1], ten conserved quantities are defined under some assumptions of asymptotic flatness, that are called Newman-Penrose (N-P) constants, but the physical meaning of these ten constants are still unclear. Much work has been done. It is found that those set of constants play important role in characterizing asymptotic flat space-time. On the level of perturbative theory, Price [2] had found the N-P constants of perturbative field strongly affected the long time evolution of that field. On the full non-linear theory, works by Friedrich, K  nn  r[3] and Valiente Kroon[4] show N-P constants contain the information that how the null infinity $\mathcal{I}^+$ touches spacial infinity i^0 . In order to get more understanding of the N-P constants, we need to consider some examples for which we can really calculate the values of these constants. People once guessed that algebraic special condition might insure zero value of them, but Kinnersley and Walker[5] showed that this idea was incorrect. Some nice works have been done to calculate the value of the N-P constants for some examples[6, 7]. In this paper, we try to see what restriction is brought by the stationary requirement. As we will see, the stationary condition and an asymptotically algebraic special condition will vanish all N-P constants.

This paper is organized as following : section II focuses on the Taylor expansion of stationary space-time. With the help of Killing equation, we reduce dynamical freedom of gravitational field into a set of free-chosen constants. Detailed expression is given up

¹e-mail : wuxn@amss.ac.cn

²e-mail : amssshangyu@yahoo.com.cn

to order $o(r^{-6})$. Section III contains the main result of this work. With algebraic special requirement, we prove all N-P constants are zero. Finally in section IV some concluding remarks are made. In Appendix, we give expression of some spin-weight harmonic functions which is useful during the proof of the main theorem.

2 The general asymptotic expansion of stationary space-time in Bondi-Sachs' coordinates

It is well-known that Bondi gauge choice[9, 10, 11] is a powerful tool to study asymptotics of space-time. In many classical works[8, 9, 10], the formal asymptotic expansion of asymptotic flat space-time has been known in Bondi coordinates. With this result, many important quantities, such as Bondi mass, Bondi energy flux, can be expressed as integral on section of null infinity. Let's focus on stationary space-times. Obviously, there is no Bondi energy flux in such space-time, i.e. $\dot{\sigma}^0 = 0$. Further more, the super-translation freedom can help us to set σ^0 to zero in stationary case [1]. Submitting this equation into the formal series, following the standard method in [9, 10], the Taylor expansions of stationary space-time are:

1) Null tetrad

$$\begin{aligned}
l^a &= \frac{\partial}{\partial r} \quad , \\
n^a &= \frac{\partial}{\partial u} + \left[-\frac{1}{2} - \frac{\Psi_2^0 + \bar{\Psi}_2^0}{2r} + \frac{\bar{\partial}\Psi_1^0 + \partial\bar{\Psi}_1^0}{6r^2} \right. \\
&\quad \left. - \frac{\bar{\partial}^2\Psi_0^0 + \partial^2\bar{\Psi}_0^0}{24r^3} - \left(\frac{|\Psi_1^0|^2}{12} + \frac{\bar{\partial}^2\Psi_0^1 + \partial^2\bar{\Psi}_0^1}{120} \right) r^{-4} + O(r^{-5}) \right] \frac{\partial}{\partial r} \\
&\quad + \left[\frac{1 + \zeta\bar{\zeta}}{6\sqrt{2}r^3} \Psi_1^0 - \frac{1 + \zeta\bar{\zeta}}{12\sqrt{2}r^4} \bar{\partial}\Psi_0^0 + O(r^{-5}) \right] \frac{\partial}{\partial \zeta} \\
&\quad + \left[\frac{1 + \zeta\bar{\zeta}}{6\sqrt{2}r^3} \bar{\Psi}_1^0 - \frac{1 + \zeta\bar{\zeta}}{12\sqrt{2}r^4} \partial\bar{\Psi}_0^0 + O(r^{-5}) \right] \frac{\partial}{\partial \bar{\zeta}} \quad , \\
m^a &= \left[-\frac{\Psi_1^0}{2r^2} + \frac{\bar{\partial}\Psi_0^0}{6r^3} + \frac{\bar{\partial}\Psi_0^1}{24r^4} + O(r^{-5}) \right] \frac{\partial}{\partial r} \\
&\quad + \left[\frac{1 + \zeta\bar{\zeta}}{6\sqrt{2}r^4} \Psi_0^0 + O(r^{-5}) \right] \frac{\partial}{\partial \zeta} + \left[\frac{1 + \zeta\bar{\zeta}}{\sqrt{2}r} + O(r^{-5}) \right] \frac{\partial}{\partial \bar{\zeta}} \quad .
\end{aligned} \tag{1}$$

2) N-P coefficients

$$\begin{aligned}
\rho &= -\frac{1}{r} + O(r^{-6}), \\
\sigma &= -\frac{\Psi_0^0}{2r^4} - \frac{\Psi_0^1}{3r^5} + O(r^{-6}), \\
\alpha &= \frac{\alpha^0}{r} - \frac{\bar{\alpha}^0\bar{\Psi}_0^0}{6r^4} + O(r^{-6}), \\
\beta &= -\frac{\bar{\alpha}^0}{r} - \frac{\Psi_1^0}{2r^3} + \frac{\alpha^0\Psi_0^0 + 2\bar{\partial}\Psi_0^0}{6r^4} + \frac{\bar{\partial}\Psi_0^1}{8r^5} + O(r^{-6}),
\end{aligned}$$

$$\begin{aligned}
\tau &= -\frac{\Psi_1^0}{2r^3} + \frac{\bar{\partial}\Psi_0^0}{3r^4} + \frac{\bar{\partial}\Psi_0^1}{8r^5} + O(r^{-6}), \\
\lambda &= -\frac{\bar{\Psi}_0^0}{12r^4} - \frac{3\bar{\Psi}_0^0\Psi_2^0 + \bar{\Psi}_0^1}{24r^5} + O(r^{-6}), \\
\mu &= -\frac{1}{2r} - \frac{\Psi_2^0}{r^2} + \frac{\bar{\partial}\Psi_1^0}{2r^3} - \frac{\bar{\partial}^2\Psi_0^0}{6r^4} - \frac{-|\Psi_1^0|^2 + \bar{\partial}^2\Psi_0^1}{6r^5} + O(r^{-6}), \\
\gamma &= -\frac{\Psi_2^0}{2r^2} + \frac{2\bar{\partial}\Psi_1^0 + \alpha^0\Psi_1^0 - \bar{\alpha}^0\bar{\Psi}_1^0}{6r^3} - \left[\frac{1}{12} (\alpha^0\bar{\partial}\Psi_0^0 - \bar{\alpha}^0\bar{\partial}\bar{\Psi}_0^0) + \frac{1}{8}\bar{\partial}^2\Psi_0^0 \right] r^{-4} \\
&\quad + \left[-\frac{1}{40} (\alpha^0\bar{\partial}\Psi_0^1 - \bar{\alpha}^0\bar{\partial}\bar{\Psi}_0^1) + \frac{1}{12}|\Psi_1^0|^2 - \frac{1}{30}\bar{\partial}^2\Psi_0^1 \right] r^{-5} + O(r^{-6}), \\
\nu &= -\left[\frac{1}{12}\bar{\Psi}_1^0 + \frac{1}{6}\bar{\partial}^2\Psi_1^0 \right] r^{-3} + \frac{1}{24} [\bar{\partial}\bar{\Psi}_0^0 + \bar{\partial}^3\Psi_0^0] r^{-4} \\
&\quad + \left[\frac{\bar{\Psi}_1^0\bar{\partial}\Psi_1^0}{20} + \frac{\Psi_2^0\bar{\partial}\bar{\Psi}_0^0}{15} + \frac{\bar{\partial}\bar{\Psi}_1^0}{80} - \frac{\bar{\Psi}_1^0\bar{\partial}\Psi_1^0}{40} - \frac{\bar{\partial}|\Psi_1^0|^2}{30} + \frac{\bar{\partial}^3\Psi_0^1}{120} \right] r^{-5} + O(r^{-6}). \quad (2)
\end{aligned}$$

3) Weyl curvature

$$\begin{aligned}
\Psi_0 &= \frac{\Psi_0^0}{r^5} + \frac{\Psi_0^1}{r^6} + O(r^{-7}), \\
\Psi_1 &= \frac{\Psi_1^0}{r^4} + \frac{\Psi_1^1}{r^5} + \frac{\Psi_1^2}{r^6} + O(r^{-7}), \\
\Psi_2 &= \frac{\Psi_2^0}{r^3} + \frac{\Psi_2^1}{r^4} + \frac{\Psi_2^2}{r^5} + \frac{\Psi_2^3}{r^6} + O(r^{-7}), \\
\Psi_3 &= \frac{\Psi_3^2}{r^4} + \frac{\Psi_3^3}{r^5} + \frac{\Psi_3^4}{r^6} + O(r^{-7}), \\
\Psi_4 &= \frac{\Psi_4^4}{r^5} + \frac{\Psi_4^5}{r^6} + O(r^{-7}).
\end{aligned} \quad (3)$$

where

$$\begin{aligned}
\Psi_1^1 &= -\bar{\partial}\Psi_0^0, \quad \Psi_1^2 = -\frac{1}{2}\bar{\partial}\Psi_0^1, \\
\Psi_2^0 &= -M, \quad \Psi_2^1 = -\frac{1}{4}\bar{\partial}\Psi_1^0, \quad \Psi_2^2 = \frac{1}{2}\bar{\partial}^2\Psi_0^0, \quad \Psi_2^3 = [-\frac{2}{3}|\Psi_1^0|^2 + \frac{1}{6}\bar{\partial}^2\Psi_0^1], \\
\Psi_3^2 &= \frac{1}{2}\bar{\partial}^2\Psi_1^0, \quad \Psi_3^3 = [-\frac{1}{2}\bar{\Psi}_1^0\Psi_2^0 - \frac{1}{6}\bar{\partial}^3\Psi_0^0], \quad \Psi_3^4 = \frac{1}{8}\bar{\Psi}_1^0\bar{\partial}\Psi_1^0 + \frac{1}{6}\bar{\partial}|\Psi_1^0|^2 - \frac{1}{24}\bar{\partial}^3\Psi_0^1, \\
\Psi_4^4 &= \frac{1}{24}\bar{\partial}^4\Psi_0^0, \quad \Psi_4^5 = \frac{9}{40}\bar{\Psi}_1^0\bar{\partial}^2\Psi_1^0 - \frac{1}{30}\bar{\partial}^3|\Psi_1^0|^2 + \frac{1}{120}\bar{\partial}^4\Psi_0^1 - \frac{1}{20}\Psi_2^0\bar{\Psi}_0^0.
\end{aligned}$$

Based on the results of characteristic initial value problem[12, 13, 14, 15], we know the dynamical freedom of asymptotic space-time are characterized by $\{\sigma^0, \Psi_1, Re(\Psi_2)\}$ on $\mathcal{I}^+$ and Ψ_4 on N , where N is an out-going light cone of the space-time. Obviously, the stationary condition has eliminated the freedom of news function. In this section, we will use Killing equation to reduce other dynamical freedom and get a general asymptotic extension of stationary space-time. As we have done in [20], we choose the standard Bondi-Sachs' coordinates, use the standard Bondi null tetrad and gauge choice[9, 10], express the time-like Killing vector as

$$t^a = Tl^a + Rn^a + \bar{A}m^a + A\bar{m}^a.$$

The Killing equations are

$$-DR = 0, \quad (4)$$

$$-DT - D'R + (\gamma + \bar{\gamma})R + \bar{\tau}A + \tau\bar{A} = 0, \quad (5)$$

$$DA - \delta R + \tau R + \bar{\rho}A + \sigma\bar{A} = 0, \quad (6)$$

$$-D'T - (\gamma + \bar{\gamma})T - \nu A - \bar{\nu}\bar{A} = 0, \quad (7)$$

$$-\tau T + \bar{\nu}R + D'A + (\bar{\gamma} - \gamma)A - \delta T - \tau T - \mu A - \bar{\lambda}\bar{A} = 0, \quad (8)$$

$$-\sigma T + \bar{\lambda}R + \delta A + (\bar{\alpha} - \beta)A = 0, \quad (9)$$

$$-\rho T + \mu R + \delta\bar{A} - (\bar{\alpha} - \beta)\bar{A} - \bar{\rho}T + \bar{\mu}R + \bar{\delta}A - (\alpha - \bar{\beta})A = 0. \quad (10)$$

Similar as the analysis in [20], we can normalize function R to 1. Assuming the asymptotic behavior of A and T as

$$\begin{aligned} T &= T^0 + \frac{T^1}{r} + \dots, \\ A &= A^0 + \frac{A^1}{r} + \dots, \end{aligned} \quad (11)$$

we can solve Killing equations order by order. The stationary condition implies $\dot{\sigma}^0 = 0$. Consequently, we have $\Psi_3^0 = 0$ and $\Psi_4^0 = 0$ because of the Bianchi identity. The first order Killing equations are

$$-A^0 = 0, \quad (12)$$

$$-\dot{T}^1 + \Psi_3^0 A^0 + \bar{\Psi}_3^0 \bar{A}^0 = 0, \quad (13)$$

$$-\bar{\Psi}_3^0 + \dot{A}^1 + \frac{1}{2}A^0 - \dot{\sigma}^0 A^0 = 0, \quad (14)$$

$$\dot{\sigma}^0 + \delta_0 A^0 + 2\bar{\alpha}^0 A^0 = 0, \quad (15)$$

$$2T^0 - 1 = 0, \quad (16)$$

which implies $\dot{T}^1 = 0$, $\dot{A}^1 = 0$.

The second order Killing equations are

$$T^1 - \frac{1}{2}(\Psi_2^0 + \bar{\Psi}_2^0) = 0, \quad (17)$$

$$-2A^1 = 0, \quad (18)$$

$$\dot{T}^2 = 0, \quad (19)$$

$$\frac{1}{2}\partial\Psi_2^0 + \dot{A}^2 - \delta_0 T^1 = 0, \quad (20)$$

$$\frac{1}{2}\sigma^0 = 0, \quad (21)$$

$$2T^1 - \sigma^0\dot{\sigma}^0 - \bar{\sigma}^0\dot{\sigma}^0 - \Psi_2^0 - \bar{\Psi}_2^0 = 0, \quad (22)$$

where $\partial f = (\delta_0 + 2s\bar{\alpha}^0)f$. From these equations, we know $\sigma^0 = 0$, $A^1 = 0$, $\dot{T}^2 = 0$, $T^1 = \frac{1}{2}(\Psi_2^0 + \bar{\Psi}_2^0)$, $\Psi_2^0 = \bar{\Psi}_2^0$, $\dot{\Psi}_2^0 = 0$, $\dot{A}^2 = -\frac{1}{2}\partial\Psi_2^0 + \frac{1}{2}\delta_0(\Psi_2^0 + \bar{\Psi}_2^0)$.

The third order equations are

$$2T^2 + \frac{1}{3}(\bar{\partial}\Psi_1^0 + \partial\bar{\Psi}_1^0) = 0, \quad (23)$$

$$-3A^2 - \frac{1}{2}\Psi_1^0 = 0, \quad (24)$$

$$\dot{T}^3 = 0, \quad (25)$$

$$\frac{1}{2}\Psi_1^0 + \bar{\nu}^3 + \dot{A}^3 + \frac{3}{2}A^2 - \delta_0 T^2 = 0, \quad (26)$$

$$\delta_0 A^2 + 2\bar{\alpha}^0 A^2 = 0, \quad (27)$$

$$2T^2 + \frac{1}{2}\bar{\partial}\Psi_1^0 + \frac{1}{2}\partial\bar{\Psi}_1^0 + \delta_0 \bar{A}^2 - 2\bar{\alpha}^0 \bar{A}^2 + \bar{\delta}_0 A^2 - 2\alpha^0 A^2 = 0. \quad (28)$$

Eq.(24),(27) imply

$$\partial\Psi_1^0 = 0. \quad (29)$$

Because the spin-weight of Ψ_1^0 is 1, we can express it as a linear combination of spin-weight harmonics, i.e.

$$\Psi_1^0 = \sum_{m=-1}^1 c_m(u) {}_1Y_{1,m}. \quad (30)$$

The action of ∂ and $\bar{\partial}$ on spin-weight harmonics are [9]

$$\begin{aligned} \partial {}_sY_{lm} &= -\sqrt{\frac{(l+s+1)(l-s)}{2}} {}_{s+1}Y_{lm}, \\ \bar{\partial} {}_sY_{lm} &= \sqrt{\frac{(l-s+1)(l+s)}{2}} {}_{s-1}Y_{lm}. \end{aligned} \quad (31)$$

Combining Eq.(23) with $\dot{T}^2 = 0$, we get

$$\begin{aligned} 0 &= \sum_{m=-1}^1 \left(\dot{c}_m {}_1Y_{1,m} + \dot{\bar{c}}_m \bar{{}_1Y_{1,m}} \right) \\ &= (\dot{c}_{-1} - \dot{\bar{c}}_1)Y_{1,-1} + (\dot{c}_0 + \dot{\bar{c}}_0)Y_{1,0} + (\dot{c}_1 - \dot{\bar{c}}_{-1})Y_{1,1}. \end{aligned}$$

This means $\dot{c}_{-1} - \dot{\bar{c}}_1 = 0$ and $\dot{c}_0 + \dot{\bar{c}}_0 = 0$. Bianchi identity $\dot{\Psi}_1^0 - \partial\Psi_2^0 = 0$ gives

$$\Psi_2^0 = -\dot{c}_{-1} {}_1Y_{1,-1} - \dot{c}_0 {}_1Y_{1,0} - \dot{c}_1 {}_1Y_{1,1} + C.$$

The stationary condition insures $\dot{\sigma}^0 = 0$, which implies $\Psi_2^0 = \bar{\Psi}_2^0$. This leads to

$$\dot{c}_{-1} + \dot{\bar{c}}_1 = 0, \quad \dot{c}_0 - \dot{\bar{c}}_0 = 0 \quad \text{and} \quad C = \bar{C}.$$

It is easy to see that $\dot{c}_m = 0$ and $\Psi_2^0 = C$, so we have

$$\begin{aligned} \Psi_1^0 &= \sum_{m=-1}^1 B_m {}_1Y_{1,m}, \\ \Psi_2^0 &= C. \end{aligned} \quad (32)$$

The Komar integral shows that $-C$ is just the Bondi mass of this space-time and $\{B_m\}$ can be understood as the components of angular momentum. Here we notice that stationary

condition helps us to restrict the form of initial data Ψ_1^0 and $Re(\Psi_2^0)$. The only freedom left is just Ψ_0 which carries information of in-going gravitational wave.

In order to get information of Ψ_0 , we need to consider higher order behavior of Killing equations. The forth order Killing equations are

$$3T^3 + (\gamma^4 + \bar{\gamma}^4) = 0, \quad (33)$$

$$4A^3 = \frac{1}{3}\bar{\partial}\Psi_0^0, \quad (34)$$

$$\dot{T}^4 + \frac{1}{3}(\bar{\partial}\Psi_1^0 + \partial\bar{\Psi}_1^0)(\Psi_2^0 + \bar{\Psi}_2^0) = 0, \quad (35)$$

$$\frac{1}{2}\Psi_1^0 T^1 - \frac{1}{3}\bar{\partial}\Psi_0^0 + \bar{\nu}^4 + \dot{A}^4 + (\Psi_2^0 + \bar{\Psi}_2^0)A^2 + \frac{3}{2}A^3 - \delta_0 T^3 + \Psi_2^0 A^2 + \frac{1}{2}A^3 = 0, \quad (36)$$

$$\frac{1}{4}\Psi_0^0 + \bar{\lambda}^4 + \partial A^3 = 0, \quad (37)$$

$$2T^3 + \mu^4 + \bar{\mu}^4 + \partial\bar{A}^3 + \bar{\partial}A^3 = 0. \quad (38)$$

Eq.(34),(37) imply

$$\Psi_0^0 = \sum_{m=-2}^2 A_m(u) {}_2Y_{2,m}, \quad (39)$$

Eq.(26),(34) insure that Ψ_0^0 is independent of u .

Our aim is to consider the Newman-Penrose constants, which are contained in coefficient Ψ_0^1 . So the fifth order of Killing equations is needed.

$$4T^4 + (\gamma^5 + \bar{\gamma}^5) - \frac{1}{2}\bar{\Psi}_1^0 A^2 - \frac{1}{2}\Psi_1^0 \bar{A}^2 = 0, \quad (40)$$

$$A^4 = \frac{1}{5}\tau^5, \quad (41)$$

$$-\frac{1}{2}\sigma^5 + \bar{\lambda}^5 + \frac{1}{2}\Psi_0^0 T^1 + \frac{3}{2}\Psi_1^0 A^2 + \partial A^4 = 0, \quad (42)$$

$$-2\rho^5 + 2T^4 + (\mu^5 + \bar{\mu}^5) + \frac{3}{2}\Psi_1^0 \bar{A}^2 + \frac{3}{2}\bar{\Psi}_1^0 A^2 + \partial\bar{A}^4 + \bar{\partial}A^4 = 0, \quad (43)$$

Eq.(41),(42) give

$$\partial\bar{\partial}\Psi_0^1 + 5\Psi_0^1 = 10(\Psi_1^0)^2 - 15\Psi_0^0\Psi_2^0. \quad (44)$$

Equation above is the key one which helps us to control the N-P constants of stationary space-time in next section. It has been noticed that each order of Killing equation helps us to control the Taylor coefficients of the same order, just like Eq.(32),(39),(44). Similarly, higher order Killing equations are also with the same property. The general form should be

$$\partial\bar{\partial}\Psi_0^k + \frac{(k+4)(k+1)}{2}\Psi_0^k = f. \quad (45)$$

where f is known function[20]. The uncertainty comes from the homogeneous part of above equations. There are some free constants in each order. These arbitrary constants should be closely related to the famous Geroch-Hansen multi-pole moments [16, 17, 18].

3 N-P constants of stationary, algebraic special space-times

In last section, we get the Taylor expansions of general asymptotic flat, stationary space-times up to order $o(r^{-6})$ which is enough for us to calculate the N-P constants. The Killing equation also helps us to restrict the dynamical freedom. We have found the information of Bondi mass is contained in Ψ_2^0 , angular momentum is in Ψ_1^0 . By definition[9], N-P constants are

$$G_m = \int_{S_\infty} {}_2Y_{2,m} \Psi_0^1 dS. \quad (46)$$

Back to eq.(44), we know

$$\not\partial \bar{\not\partial} \Psi_0^1 + 5\Psi_0^1 = 10(\Psi_1^0)^2 - 15\Psi_0^0 \Psi_2^0. \quad (47)$$

This is an inhomogeneous linear equation. The associated homogeneous equation is

$$\not\partial \bar{\not\partial} \Psi + 5\Psi = 0. \quad (48)$$

Keeping Eq.(31) in mind, we can get the general solution of above equation:

$$\Psi = \sum_{m=-3}^3 D_m {}_2Y_{3,m} \quad .$$

Let $3\tilde{\Psi}_0^1 = 10(\Psi_1^0)^2 - 15\Psi_2^0 \Psi_0^0$, with the help of Eq.(32) and (39), we know

$$\not\partial \bar{\not\partial} \tilde{\Psi}_0^1 + 5\tilde{\Psi}_0^1 = 10(\Psi_1^0)^2 - 15\Psi_0^0 \Psi_2^0, \quad (49)$$

then the general solution of Eq.(47) is

$$\Psi_0^1 = \tilde{\Psi}_0^1 + \sum_{m=-3}^3 D_m {}_2Y_{3,m} \quad . \quad (50)$$

Because of the definition, it is clear that only $\tilde{\Psi}_0^1$ contributes to the N-P constants. Combining above equation with Eq.(32),(39), we have

$$\begin{aligned} \tilde{\Psi}_0^1 &= \frac{10}{3}(\Psi_1^0)^2 - 5\Psi_2^0 \Psi_0^0 \\ &= \frac{5}{3} \times [2(\sum_{m=-1}^1 B_m {}_1Y_{1,m})^2 + 3M(\sum_{m=-2}^2 A_m {}_2Y_{2,m})] \\ &= \frac{10}{3}[B_{-1} {}_1Y_{1,-1} + B_0 {}_1Y_{1,0} + B_1 {}_1Y_{1,1}]^2 \\ &\quad + 5M[A_{-2} {}_2Y_{2,-2} + A_{-1} {}_2Y_{2,-1} + A_0 {}_2Y_{2,0} + A_1 {}_2Y_{2,-1} + A_2 {}_2Y_{2,2}] \\ &= \frac{10}{3}[(B_{-1})^2({}_1Y_{1,-1})^2 + (B_0)^2({}_1Y_{1,0})^2 + (B_1)^2({}_1Y_{1,1})^2 \\ &\quad + 2B_{-1}B_0 {}_1Y_{1,-1} {}_1Y_{1,0} + 2B_0B_1 {}_1Y_{1,0} {}_1Y_{1,1} + B_{-1}B_1 {}_1Y_{1,-1} {}_1Y_{1,1}] \\ &\quad + 5M[A_{-2} {}_2Y_{2,-2} + A_{-1} {}_2Y_{2,-1} + A_0 {}_2Y_{2,0} + A_1 {}_2Y_{2,-1} + A_2 {}_2Y_{2,2}] \end{aligned}$$

$$\begin{aligned}
&= \frac{5}{3} \left\{ [2(B_{-1})^2 \sqrt{\frac{3}{40\pi}} + 3MA_{-2}] {}_2Y_{2,-2} + [4B_{-1}B_0 \sqrt{\frac{3}{80\pi}} + 3MA_{-1}] {}_2Y_{2,-1} \right. \\
&\quad + [4(B_0)^2 \sqrt{\frac{1}{80\pi}} + 4B_{-1}B_1 \sqrt{\frac{1}{80\pi}} + 3MA_0] {}_2Y_{2,0} \\
&\quad \left. + [4B_0B_1 \sqrt{\frac{3}{80\pi}} + 3MA_1] {}_2Y_{2,1} + [2(B_1)^2 \sqrt{\frac{3}{40\pi}} + 3MA_2] {}_2Y_{2,2} \right\} \\
&= \sum_{m=-2}^2 G_m {}_2Y_{2,m}.
\end{aligned} \tag{51}$$

By definition, $\{G_m\}$ are just N-P constants. In general, it's easy to understand that N-P constants may not be zero because there are free constants $\{B_m\}$ and $\{A_m\}$ which are out of our control.

Now, we consider the algebraic special case. The algebraic special condition is [19]

$$\begin{aligned}
I^3 &= 27J^2, \\
I &= \Psi_0\Psi_4 - 4\Psi_1\Psi_3 + 3(\Psi_2)^2, \\
J &= \Psi_4\Psi_2\Psi_0 + 2\Psi_3\Psi_2\Psi_1 - (\Psi_2)^3 - (\Psi_3)^2\Psi_0 - (\Psi_1)^2\Psi_4.
\end{aligned} \tag{52}$$

Submitting the asymptotic extension of Weyl curvature into above equations, we get the first non-zero equation as

$$\begin{aligned}
0 &= 36(\Psi_1^0)^2(\Psi_2^0)^2(\Psi_3^0)^2 - 54(\Psi_2^0)^3(\Psi_3^0)^2\Psi_0^0 - 54(\Psi_1^0)^2(\Psi_2^0)^3\Psi_4^0 + 81(\Psi_2^0)^4\Psi_0^0\Psi_4^0 \\
&= 9[(\Psi_1^0)^2(\bar{\partial}^2\Psi_1^0)^2 + \frac{3M}{2}(\bar{\partial}^2\Psi_1^0)^2\Psi_0^0 + 6M(\Psi_1^0)^2\frac{1}{24}\bar{\partial}^4\Psi_0^0 + \frac{9}{24}M^2\Psi_0^0\bar{\partial}^4\Psi_0^0] \\
&= 9[(\sum_{m=-1}^1 B_{m-1}Y_{1,m})^2(\sum_{m=-1}^1 B_{m-1}Y_{1,m})^2 + \frac{3M}{2}(\sum_{m=-1}^1 B_{m-1}Y_{1,m})^2(\sum_{m=-2}^2 A_{m-2}Y_{2,m}) \\
&\quad + \frac{3M}{2}(\sum_{m=-1}^1 B_{m-1}Y_{1,m})^2(\sum_{m=-2}^2 A_{m-2}Y_{2,m}) + \frac{9M^2}{4}(\sum_{m=-2}^2 A_{m-2}Y_{2,m})(\sum_{m=-2}^2 A_{m-2}Y_{2,m})] \\
&= \frac{9}{4} \times [2(\sum_{m=-1}^1 B_{m-1}Y_{1,m})^2 + 3M(\sum_{m=-2}^2 A_{m-2}Y_{2,m})] \\
&\quad \times [2(\sum_{m=-1}^1 B_{m-1}Y_{1,m})^2 + 3M(\sum_{m=-2}^2 A_{m-2}Y_{2,m})],
\end{aligned} \tag{53}$$

which means

$$2(\sum_{m=-1}^1 B_{m-1}Y_{1,m})^2 + 3M(\sum_{m=-2}^2 A_{m-2}Y_{2,m}) = 0 \tag{54}$$

or

$$2(\sum_{m=-1}^1 B_{m-1}Y_{1,m})^2 + 3M(\sum_{m=-2}^2 A_{m-2}Y_{2,m}) = 0. \tag{55}$$

Comparing Eq.(54) with the second line of eq.(51), it is easy to see that N-P constants vanish if Eq.(54) holds. If Eq.(55) holds, we have

$$0 = 2(\sum_{m=-1}^1 B_{m-1}Y_{1,m})^2 + 3M(\sum_{m=-2}^2 A_{m-2}Y_{2,m})$$

$$\begin{aligned}
&= 2[B_1^2(-{}_1Y_{1,1})^2 + B_0^2(-{}_1Y_{1,0})^2 + B_{-1}^2(-{}_1Y_{1,-1})^2 \\
&\quad + 2B_1B_0{}_1Y_{1,1}{}_1Y_{1,0} + 2B_1B_{-1}{}_1Y_{1,1}{}_1Y_{1,-1} + 2B_0B_{-1}{}_1Y_{1,0}{}_1Y_{1,-1}] \\
&\quad + 3MA_2{}_2Y_{2,2} + 3MA_1{}_2Y_{2,1} + 3MA_0{}_2Y_{2,0} + 3MA_{-1}{}_2Y_{2,-1} + 3MA_{-2}{}_2Y_{2,-2} \\
&= \left[2\sqrt{\frac{3}{40\pi}}B_1^2 + 3MA_2\right]{}_2Y_{2,2} + \left[4\sqrt{\frac{3}{80\pi}}B_1B_0 + 3MA_1\right]{}_2Y_{2,1} \\
&\quad + \left[\frac{4}{\sqrt{80\pi}}(B_0^2 + B_1B_{-1}) + 3MA_0\right]{}_2Y_{2,0} \\
&\quad + \left[4B_{-1}B_0\sqrt{\frac{3}{80\pi}} + 3MA_{-1}\right]{}_2Y_{2,-1} + \left[2(B_{-1})^2\sqrt{\frac{3}{40\pi}} + 3MA_{-2}\right]{}_2Y_{2,-2}. \quad (56)
\end{aligned}$$

Comparing above equation with the fifth line of Eq.(51), Eq.(55) also implies zero value of the N-P constants, so we have proved following result

Theorem 1 *All N-P constants of asymptotic flat, stationary, algebraic special space-time vanish.*

Remark : What needs to emphasis is that Kerr space-time is an important special case of our theorem, i.e. all N-P constants of Kerr space-time are zero. Based on our knowledge, the value of the N-P constants of Kerr space-time is unknown. The value of N-P constants of Kerr space-time are also got by us with other methods[20, 21].

4 Discussion

In last section, we prove a theorem that asymptotic flat, stationary, algebraic special condition vanishes all N-P constants. In fact, our result is a bit stronger than above theorem because only the leading order of the algebraic special condition is needed. Maybe we can call such kind of space-time as asymptotically algebraic special space-time. This tells us that algebraic special condition may not be a very suitable choice to control the N-P constants. A natural question is whether the stationary condition is enough to vanish all N-P constants in vacuum case. As we have seen, the stationary condition has strongly restricted the freedom of gravitational field. If this idea is right, it would give a physical understanding for N-P constants. Although we still haven't very strong evidence, some primary calculation implies this idea maybe right. This will be discussed in future works.

Acknowledgement

This work is supported by National Science Foundation of China under Grant No. 10375087 and K.C.Wong Education Foundation, Hong Kong. Authors would like to thank Prof.Y.K.Lau, Dr.Z.J.Cao, S.Bai and X.F.Gong for their helpful discussions. X.Wu also wants to thank Prof.H.Friedrich and Dr.J.A.Valiente Kroon for their helpful discussion.

Appendix

Some spin-weight harmonics

$$\begin{aligned}
Y_{0,0} &= \frac{1}{\sqrt{4\pi}}; \\
{}_1Y_{1,1} &= \sqrt{\frac{3}{16\pi}}(\cos\theta + 1)e^{i\phi}, \\
{}_1Y_{1,0} &= \sqrt{\frac{3}{8\pi}}\sin\theta, \\
{}_1Y_{1,-1} &= \sqrt{\frac{3}{16\pi}}(1 - \cos\theta)e^{-i\phi}; \\
Y_{1,1} &= -\sqrt{\frac{3}{8\pi}}\sin\theta e^{i\phi}, \\
Y_{1,0} &= \sqrt{\frac{3}{4\pi}}\cos\theta, \\
Y_{1,-1} &= \sqrt{\frac{3}{8\pi}}\sin\theta e^{-i\phi}; \\
{}_{-1}Y_{1,1} &= \sqrt{\frac{3}{16\pi}}(1 - \cos\theta)e^{i\phi}, \\
{}_{-1}Y_{1,0} &= -\sqrt{\frac{3}{8\pi}}\sin\theta, \\
{}_{-1}Y_{1,-1} &= \sqrt{\frac{3}{16\pi}}(1 + \cos\theta)e^{-i\phi}; \\
{}_2Y_{2,2} &= 3\sqrt{\frac{5}{96\pi}}(1 + \cos\theta)^2e^{2i\phi}, \\
{}_2Y_{2,1} &= 3\sqrt{\frac{5}{24\pi}}\sin\theta(1 + \cos\theta)e^{i\phi}, \\
{}_2Y_{2,0} &= 3\sqrt{\frac{5}{16\pi}}\sin^2\theta, \\
{}_2Y_{2,-1} &= 3\sqrt{\frac{5}{24\pi}}\sin\theta(1 - \cos\theta)e^{-i\phi}, \\
{}_2Y_{2,-2} &= 3\sqrt{\frac{5}{96\pi}}(1 - \cos\theta)^2e^{-2i\phi}; \\
{}_1Y_{2,2} &= -3\sqrt{\frac{5}{24\pi}}\sin\theta(1 + \cos\theta)e^{2i\phi}, \\
{}_1Y_{2,1} &= 3\sqrt{\frac{5}{24\pi}}(2\cos\theta - 1)(1 + \cos\theta)e^{i\phi}, \\
{}_1Y_{2,0} &= 3\sqrt{\frac{5}{4\pi}}\sin\theta\cos\theta,
\end{aligned}$$

$$\begin{aligned}
{}_1Y_{2,-1} &= 3\sqrt{\frac{5}{24\pi}}(2\cos\theta+1)(1-\cos\theta)e^{-i\phi}, \\
{}_1Y_{2,-2} &= 3\sqrt{\frac{5}{24\pi}}\sin\theta(1-\cos\theta)e^{-2i\phi}; \\
Y_{2,2} &= 3\sqrt{\frac{5}{16\pi}}\sin^2\theta e^{2i\phi}, \\
Y_{2,1} &= -6\sqrt{\frac{5}{16\pi}}\sin\theta\cos\theta e^{i\phi}, \\
Y_{2,0} &= \sqrt{\frac{5}{24\pi}}(3\cos^2\theta-1), \\
Y_{2,-1} &= 6\sqrt{\frac{5}{16\pi}}\sin\theta\cos\theta e^{-i\phi}, \\
Y_{2,-2} &= 3\sqrt{\frac{5}{16\pi}}\sin^2\theta e^{-2i\phi}; \\
{}_{-1}Y_{2,2} &= -3\sqrt{\frac{5}{24\pi}}\sin\theta(1-\cos\theta)e^{2i\phi}, \\
{}_{-1}Y_{2,1} &= 3\sqrt{\frac{5}{24\pi}}(2\cos\theta+1)(1-\cos\theta)e^{i\phi}, \\
{}_{-1}Y_{2,0} &= -\sqrt{\frac{5}{4\pi}}\sin\theta\cos\theta, \\
{}_{-1}Y_{2,-1} &= 3\sqrt{\frac{5}{24\pi}}(2\cos\theta-1)(1+\cos\theta)e^{-i\phi}, \\
{}_{-1}Y_{2,-2} &= 3\sqrt{\frac{5}{24\pi}}\sin\theta(1+\cos\theta)e^{-2i\phi}; \\
{}_{-2}Y_{2,2} &= 3\sqrt{\frac{5}{96\pi}}(1-\cos\theta)^2e^{2i\phi}, \\
{}_{-2}Y_{2,1} &= -3\sqrt{\frac{5}{24\pi}}\sin\theta(1-\cos\theta)e^{i\phi}, \\
{}_{-2}Y_{2,0} &= 3\sqrt{\frac{5}{16\pi}}\sin^2\theta, \\
{}_{-2}Y_{2,-1} &= -3\sqrt{\frac{5}{24\pi}}\sin\theta(1+\cos\theta)e^{-i\phi}, \\
{}_{-2}Y_{2,-2} &= 3\sqrt{\frac{5}{96\pi}}(1+\cos\theta)^2e^{-2i\phi}.
\end{aligned}$$

References

- [1] E. T. Newman and R. Penrose, Proc. Roy. Soc. Lond. **A 305** (1968) 175.
- [2] R. H. Price, Phys. Rev. **D 5** (1972) 2419.

- [3] H. Friedrich and J. Kánnár, J. Math. Phys. **41** (2000) 2195.
- [4] J.A. Valiente Kroon, Class. Quant. Grav. **20** (2003) L53.
- [5] W. Kinnersely and M. Walker, Phys. Rev. **D 2** (1970) 1359.
- [6] R. Lazkoz and J. A. Valiente Kroon, Phys. Rev. **D62** (2000) 084033.
- [7] S. Dain and J. A. Valiente Kroon, Class. Quant. Grav., **19**(2002)811.
- [8] H. Bondi, M. G. J. van der Burg and A.W.K.Metzner, Proc. Roy. Soc. Lond. **A 269** (1962) 21.
- [9] R. Penrose and R. Rindler, *Spinors and Space-Time* Vol.I and II, Cambridge University Press, 1986.
- [10] E. T. Newman and T. W. J. Unti, J. Math. Phys. **3** (1962) 891.
- [11] J. Stewart, *Advanced General Relativity*, Cambridge University Press, 1990.
- [12] H. Friedrich, Proc. Roy. Soc. Lond. **A 378** (1981) 169-184, 401-421
- [13] H. Friedrich, Proc. Roy. Soc. Lond. **A 381** 361-371.
- [14] J. Kánnár, Proc. Roy. Soc. Lond. **A 452** (1996) 945.
- [15] F. Nicoló, “A local characteristic problem for the Einstein vacuum equations”, gr-qc/0603118.
- [16] R. Geroch, J. Math. Phys. **11** (1970) 1955, 2580.
- [17] R. Hansen, J. Math. Phys. **15** (1974) 46.
- [18] H. Friedrich, “Static vacuum solutions from convergent null data expansions at space-like infinity”, gr-qc/0606133.
- [19] D. Kramer, H. Stephani, E. Herlt and M. MacCallum, *Exact Solutions of Einstein’s Field Equations*, Cambridge University Press, 1980.
- [20] X. Wu and S. Bai, “On local uniqueness of Kerr space-time”, in preparing.
- [21] S. Bai et.al., “Bondi-Sachs formulism of Kerr metric”, submit to Phys. Rev. **D**.